

18/20

Figure 18

SEQ ID NO:1




Homo Sapiens Ephrin type-B receptor 4 Precursor (EphB4)

1	MELRVLLCWA	SLAAALEETL	LNTKLETADL	KWVTFPQVDG	QWEELSGLDE	EQHSVRITYEV
61	CDVQRAPGQA	HWLRTGWVPR	RGAVHVIYATL	RFTMLECLSL	PRAGRSCKET	FTVFYYESDA
21	DTATALTPAW	MENPYIKVDT	VAAEHLTRKR	PGAEATGKVN	VKTLRLGPLS	KAGFYLAFAQD
181	QGACMALLSL	HLFYKKCAQL	TVNLTRFPET	VPRELVVPVA	GSCVVDVPA	PGPSPSLYCR
241	EDGQWAEQPV	TGCSCAPGFE	AAEGNTKCRA	CAQGTFKPLS	GEGSCQPCPA	NSHSNTIGSA
301	VCQCRVGYFR	ARTDPRGAPC	TTPPSAPRSV	VSRLNGSSLH	LEWSAPLESG	GREDLTYALR
361	CRECRPGGSC	APCGGDLTFD	PGPRDLVEPW	VVVRGLRPDF	TYTFEVTALN	GVSSLATGPV
421	PFEPVNVTTD	REVPPAVSDI	RVTRSSPSSL	SLAWAVPRAP	SGAVLDYEVK	YHEKGAEGPS
481	SVRFLKTSN	RAELRGLKRG	ASYLVQVRAR	SEAGYGPFQ	EHHSQTQLDE	SEGWREQLAL
541	IAGTAVVGVV	LVLVVIVVAV	LCLRKQSNR	EAEYSDKHGQ	YLIGHGIKVY	IDPFTYEDPN
601	EAVREFAKEI	DVSIVKIEEV	IGAGEFGEVC	RGRLKAPGKK	ESCVAIKTLK	GGYTERQRRE
661	FLSEASIMGQ	FEHPNIIRLE	GVVTNSMPVM	ILTEFMENGA	LDSFLRLNDG	QFTVIQLVGM
721	LRGIASGMRY	LAEMSYVHRD	LAARNILVNS	NLVCKVSDFG	LSRFLEENSS	DPTYTSSLGG
781	KIPRWTAPE	AIAFRKFTSA	SDAWSYGIVM	WEVMSFGERP	YWDMSNQDVI	NAIEQDYRLP
841	PPDCPTSLH	QLMLDCWQKD	RNARPRFPQV	VSALDKMIRN	PASLKIVARE	NGGASHPLLD
901	QRQPHYSAFG	SVGWLRAIK	MGRYEEFAA	AGFGSFELVS	QISAEDLLRI	GVTLAGHQKK
961	ILASVQHMK	QAKPGTPGGT	GGPAPQY			

Added Q

Added Y

Added N

   My NCBI [\[Sign In\]](#) [\[Regis\]](#)

PubMed Nucleotide Protein Genome Structure PMC Taxonomy OMIM Books

Search for

Display Show Send to

Range: from to Features: version 2

☐ 1: NP_004435. Reports ...[gi:17975770] The record has been replaced by NP_004435.3

Comment Features Sequence

LOCUS NP_004435 987 aa linear PRI 05-APR-2003
DEFINITION ephrin receptor EphB4 precursor; Ephrin receptor EphB4 (hepatoma transmembrane kinase); Tyroll; ephrin receptor EphB4; hepatoma transmembrane kinase [Homo sapiens].
ACCESSION NP_004435
VERSION NP_004435.2 GI:17975770
DBSOURCE REFSEQ: accession NM_004444.2
KEYWORDS .
SOURCE Homo sapiens (human)
ORGANISM Homo sapiens
Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi; Mammalia; Eutheria; Primates; Catarrhini; Hominidae; Homo.
REFERENCE 1 (residues 1 to 987)
AUTHORS Steinle,J.J., Meininger,C.J., Forough,R., Wu,G., Wu,M.H. and Granger,H.J.
TITLE Eph B4 receptor signaling mediates endothelial cell migration and proliferation via the phosphatidylinositol 3-kinase pathway
JOURNAL J. Biol. Chem. 277 (46), 43830-43835 (2002)
PUBMED 12235151
REMARK GeneRIF: Eph B4 stimulates migration and proliferation and may play a role in angiogenesis
REFERENCE 2 (residues 1 to 987)
AUTHORS Suenobu,S., Takakura,N., Inada,T., Yamada,Y., Yuasa,H., Zhang,X.Q., Sakano,S., Oike,Y. and Suda,T.
TITLE A role of EphB4 receptor and its ligand, ephrin-B2, in erythropoiesis
JOURNAL Biochem. Biophys. Res. Commun. 293 (3), 1124-1131 (2002)
PUBMED 12051776
REMARK GeneRIF: role of EphB4 receptor in erythropoiesis
REFERENCE 3 (residues 1 to 987)
AUTHORS Wang,Z., Miura,N., Bonelli,A., Mole,P., Carlesso,N., Olson,D.P. and Scadden,D.T.
TITLE Receptor tyrosine kinase, EphB4 (HTK), accelerates differentiation of select human hematopoietic cells
JOURNAL Blood 99 (8), 2740-2747 (2002)
PUBMED 11929761
REMARK GeneRIF: accelerates differentiation of select human hematopoietic cells
REFERENCE 4 (residues 1 to 987)
AUTHORS Wilson,M.D., Riemer,C., Martindale,D.W., Schnupf,P., Boright,A.P., Cheung,T.L., Hardy,D.M., Schwartz,S., Scherer,S.W., Tsui,L.C., Miller,W. and Koop,B.F.
TITLE Comparative analysis of the gene-dense ACHE/TFR2 region on human

chromosome 7q22 with the orthologous region on mouse chromosome 5

JOURNAL Nucleic Acids Res. 29 (6), 1352-1365 (2001)
PUBMED [11239002](#)

REFERENCE 5 (residues 1 to 987)

AUTHORS Berclaz,G., Andres,A.C., Albrecht,D., Dreher,E., Ziemiecki,A.,
Gusterson,B.A. and Crompton,M.R.

TITLE Expression of the receptor protein tyrosine kinase myk-1/htk in
normal and malignant mammary epithelium

JOURNAL Biochem. Biophys. Res. Commun. 226 (3), 869-875 (1996)
PUBMED [8831703](#)

REFERENCE 6 (residues 1 to 987)

AUTHORS Bennett,B.D., Wang,Z., Kuang,W.J., Wang,A., Groopman,J.E.,
Goeddel,D.V. and Scadden,D.T.

TITLE Cloning and characterization of HTK, a novel transmembrane tyrosine
kinase of the EPH subfamily

JOURNAL J. Biol. Chem. 269 (19), 14211-14218 (1994)
PUBMED [8188704](#)

REFERENCE 7 (residues 1 to 987)

AUTHORS Andres,A.C., Reid,H.H., Zurcher,G., Blaschke,R.J., Albrecht,D. and
Ziemiecki,A.

TITLE Expression of two novel eph-related receptor protein tyrosine
kinases in mammary gland development and carcinogenesis

JOURNAL Oncogene 9 (5), 1461-1467 (1994)
PUBMED [8152808](#)

COMMENT REVIEWED REFSEQ: This record has been curated by NCBI staff. The
reference sequence was derived from [U07695.1](#).
[WARNING] On Jul 11, 2003 this sequence was replaced by
gi:[32528301](#).
→ On Dec 21, 2001 this sequence version replaced gi:[4758290](#).

Summary: Ephrin receptors and their ligands, the ephrins, mediate numerous developmental processes, particularly in the nervous system. Based on their structures and sequence relationships, ephrins are divided into the ephrin-A (EFNA) class, which are anchored to the membrane by a glycosylphosphatidylinositol linkage, and the ephrin-B (EFNB) class, which are transmembrane proteins. The Eph family of receptors are divided into 2 groups based on the similarity of their extracellular domain sequences and their affinities for binding ephrin-A and ephrin-B ligands. Ephrin receptors make up the largest subgroup of the receptor tyrosine kinase (RTK) family. The protein encoded by this gene binds to ephrin-B2 and plays an essential role in vascular development.

FEATURES Location/Qualifiers

source 1..987
/organism="Homo sapiens"
/db_xref="taxon:[9606](#)"
/chromosome="7"
/map="7q22"

Protein 1..987
/product="ephrin receptor EphB4 precursor"
/EC_number="[2.7.1.112](#)"
/note="Ephrin receptor EphB4 (hepatoma transmembrane kinase); Tyro11; ephrin receptor EphB4"

sig_peptide 1..15
/calculated_mol_wt=1647

mat_peptide 16..987
/product="ephrin receptor EphB4"
/calculated_mol_wt=106743

Region 29..197
/region_name="Ephrin receptor ligand binding domain. The

Eph receptors, which bind to ephrins pfam00812 are a large family of receptor tyrosine kinases. This family represents the amino terminal domain which binds the ephrin ligand"

/note="EPH_lbd; Ephrin receptor ligand binding domain. The Eph receptors, which bind to ephrins pfam00812 are a large family of receptor tyrosine kinases. This family represents the amino terminal domain which binds the ephrin ligand"

/db_xref="CDD:pfam01404"

Region 255..313

/region_name="Giardia variant-specific surface protein"

/note="VSP; Giardia variant-specific surface protein"

/db_xref="CDD:pfam03302"

Region 324..414

/region_name="Fibronectin type 3 domain"

/note="FN3; Fibronectin type 3 domain"

/db_xref="CDD:smart00060"

Region 437..517

/region_name="Fibronectin type III domain"

/note="fn3; Fibronectin type III domain"

/db_xref="CDD:pfam00041"

Region 615..874

/region_name="Tyrosine kinase, catalytic domain"

/note="TyrKc; Tyrosine kinase, catalytic domain"

/db_xref="CDD:smart00219"

Region 904..971

/region_name="Sterile alpha motif. Widespread domain in signalling and nuclear proteins. In EPH-related tyrosine kinases, appears to mediate cell-cell initiated signal transduction via the binding of SH2-containing proteins to a conserved tyrosine that is phosphorylated. In many cases mediates homodimerisation"

/note="SAM; Sterile alpha motif. Widespread domain in signalling and nuclear proteins. In EPH-related tyrosine kinases, appears to mediate cell-cell initiated signal transduction via the binding of SH2-containing proteins to a conserved tyrosine that is phosphorylated. In many cases mediates homodimerisation"

/db_xref="CDD:smart00454"

CDS 1..987

/gene="EPHB4"

/coded_by="NM_004444.2:86..3049"

/db_xref="GeneID:2050"

/db_xref="MIM:600011"

ORIGIN

```

1 melrvllcwa slaaaleetl lntkletadl kwvtfpqvdg qweelsglde eqhsvrtyev
61 cevqrapgqa hwlrtgwvpr rgavhvayat rftmleclsl pragrcket ftvfyyesda
121 dtataltpaw menpyikvdt vaaehltrkr pgaeatgkvn vktlrlgpls kagfylaifqd
181 qgacmallsl hlffykcqal tvnltrfpet vprelvvpva gscvvdavpa pgpspslycr
241 edgqwaeqp v tgcscapgfe aaegntkcra caqgtfkpls gegscqpcpa nshsntigsa
301 vcqcrvgyfr artdprgapc ttpsaprsv vsrlngsslh lewsaplesg gredltyalr
361 crecrpggsc apcggdltfd pgprdlvepw vvvrgrlpdf tytftevtaln gvsslatgvp
421 pfepvnvtt d revppavsd i rvtrsspssl slawavprap sgawldyevk yhekgaegps
481 svrflktsen raelrglkr g asylvqvrar seagygpfgq ehhsqtqlde segwreqlal
541 iagtavvgvv lylvvivvav lclrkqsngr eaeysdkhgq ylighgtkvy idpftyedpn
601 eavrefakei dvsyvkieev igagefgevc rgrlkapgk escvaiktlk ggyterqrre
661 flseasimgq fehpnirle gvvtnsmpvm iltefmenga ldsflrlndg qftviqlvgm
721 lrgiasgmry laemsvhrd laarnilvns nlvckvsdfg lsrflenss dptytsslgg
781 kipirwtape aiafrkftsa sdawsygivm wevmsfgerp ywdmsnqdv naieqdyrlp

```



841 pppdcptslh qlmldcwqkd rnarprfpqv vsalckmirn paslkivare nggashplld
901 qrqphysafg svgewlraik mgryeesfaa agfgsfelvs qisaedllri gvtlaghqkk
961 ilasvqhmks qakpgtpggt ggpapqy

//

[Disclaimer](#) | [Write to the Help Desk](#)
[NCBI](#) | [NLM](#) | [NIH](#)

Last update: Tue, 29 Apr 2008 Rev. 126150

Exhibit B

  My NCBI [Sign In] [Regis]

PubMed Nucleotide Protein Genome Structure PMC Taxonomy OMIM Books

Search for

Display Show Send to

Range: from to Features: ☐ SNP ☒ CDD Version 3

☐ 1: NP_004435. Reports ephrin receptor E...[gi:32528301]BLink, Conserved
Domains, LinksComment Features Sequence

LOCUS NP_004435 987 aa linear PRI 29-APR-2008
DEFINITION ephrin receptor EphB4 precursor [Homo sapiens].
ACCESSION NP_004435
VERSION NP_004435.3 GI:32528301
DBSOURCE REFSEQ: accession NM_004444.4
KEYWORDS .
SOURCE Homo sapiens (human)
ORGANISM Homo sapiens
Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi;
Mammalia; Eutheria; Euarchontoglires; Primates; Haplorrhini;
Catarrhini; Hominidae; Homo.
REFERENCE 1 (residues 1 to 987)
AUTHORS Alam,S.M., Fujimoto,J., Jahan,I., Sato,E. and Tamaya,T.
TITLE Coexpression of EphB4 and ephrinB2 in tumour advancement of ovarian
cancers
JOURNAL Br. J. Cancer 98 (4), 845-851 (2008)
PUBMED 18231102
REMARK GeneRIF: Overexpression of EphB4 is associated with tumour
advancement of ovarian cancers
REFERENCE 2 (residues 1 to 987)
AUTHORS Tachibana,M., Tonomoto,Y., Hyakudomi,R., Hyakudomi,M., Hattori,S.,
Ueda,S., Kinugasa,S. and Yoshimura,H.
TITLE Expression and prognostic significance of EFNB2 and EphB4 genes in
patients with oesophageal squamous cell carcinoma
JOURNAL Dig Liver Dis 39 (8), 725-732 (2007)
PUBMED 17611172
REMARK GeneRIF: EphB4 expression was found in 44 (72.1%) out of 61 cancer
tissues analysed.
REFERENCE 3 (residues 1 to 987)
AUTHORS Foubert,P., Silvestre,J.S., Souttou,B., Barateau,V., Martin,C.,
Ebrahimian,T.G., Lere-Dean,C., Contreres,J.O., Sulpice,E.,
Levy,B.I., Plouet,J., Tobelem,G. and Le Ricousse-Roussanne,S.
TITLE PSGL-1-mediated activation of EphB4 increases the proangiogenic
potential of endothelial progenitor cells
JOURNAL J. Clin. Invest. 117 (6), 1527-1537 (2007)
PUBMED 17510705
REMARK GeneRIF: activation of EphB4 enhances proangiogenic capacity
through induction of PSGL-1 expression and adhesion to E selectin
and P selectin
REFERENCE 4 (residues 1 to 987)
AUTHORS Kumar,S.R., Masood,R., Spannuth,W.A., Singh,J., Scehnet,J.,
Kleiber,G., Jennings,N., Deavers,M., Krasnoperov,V., Dubeau,L.,
Weaver,F.A., Sood,A.K. and Gill,P.S.
TITLE The receptor tyrosine kinase EphB4 is overexpressed in ovarian

cancer, provides survival signals and predicts poor outcome

JOURNAL Br. J. Cancer 96 (7), 1083-1091 (2007)

PUBMED [17353927](#)

REMARK GeneRIF: Overexpression of EphB4 is associated with ovarian cancer

REFERENCE 5 (residues 1 to 987)

AUTHORS Limbourg,A., Ploom,M., Elligsen,D., Sorensen,I., Ziegelhoeffer,T., Gossler,A., Drexler,H. and Limbourg,F.P.

TITLE Notch ligand Delta-like 1 is essential for postnatal arteriogenesis

JOURNAL Circ. Res. 100 (3), 363-371 (2007)

PUBMED [17234965](#)

REMARK GeneRIF: Notch signaling by induction of Dll1 was necessary & sufficient to induce EphB4-dependent branching morphogenesis in human arterial EC.

REFERENCE 6 (residues 1 to 987)

AUTHORS Zhou,R.

TITLE The Eph family receptors and ligands

JOURNAL Pharmacol. Ther. 77 (3), 151-181 (1998)

PUBMED [9576626](#)

REMARK Review article

REFERENCE 7 (residues 1 to 987)

AUTHORS Flanagan,J.G. and Vanderhaeghen,P.

TITLE The ephrins and Eph receptors in neural development

JOURNAL Annu. Rev. Neurosci. 21, 309-345 (1998)

PUBMED [9530499](#)

REMARK Review article

REFERENCE 8 (residues 1 to 987)

AUTHORS Berclaz,G., Andres,A.C., Albrecht,D., Dreher,E., Ziemiecki,A., Gusterson,B.A. and Crompton,M.R.

TITLE Expression of the receptor protein tyrosine kinase myk-1/htk in normal and malignant mammary epithelium

JOURNAL Biochem. Biophys. Res. Commun. 226 (3), 869-875 (1996)

PUBMED [8831703](#)

REFERENCE 9 (residues 1 to 987)

AUTHORS Bennett,B.D., Wang,Z., Kuang,W.J., Wang,A., Groopman,J.E., Goeddel,D.V. and Scadden,D.T.

TITLE Cloning and characterization of HTK, a novel transmembrane tyrosine kinase of the EPH subfamily

JOURNAL J. Biol. Chem. 269 (19), 14211-14218 (1994)

PUBMED [8188704](#)

REFERENCE 10 (residues 1 to 987)

AUTHORS Andres,A.C., Reid,H.H., Zurcher,G., Blaschke,R.J., Albrecht,D. and Ziemiecki,A.

TITLE Expression of two novel eph-related receptor protein tyrosine kinases in mammary gland development and carcinogenesis

JOURNAL Oncogene 9 (5), 1461-1467 (1994)

PUBMED [8152808](#)

REMARK Erratum:[Oncogene 1994 Aug;9(8):2431]

COMMENT REVIEWED REFSEQ: This record has been curated by NCBI staff. The reference sequence was derived from [CB962032.1](#), [BC052804.1](#) and [AY056048.1](#).



On Jul 11, 2003 this sequence version replaced gi:[17975770](#).

Summary: Ephrin receptors and their ligands, the ephrins, mediate numerous developmental processes, particularly in the nervous system. Based on their structures and sequence relationships, ephrins are divided into the ephrin-A (EFNA) class, which are anchored to the membrane by a glycosylphosphatidylinositol linkage, and the ephrin-B (EFNB) class, which are transmembrane proteins. The Eph family of receptors are divided into 2 groups based on the similarity of their extracellular domain sequences and their

affinities for binding ephrin-A and ephrin-B ligands. Ephrin receptors make up the largest subgroup of the receptor tyrosine kinase (RTK) family. The protein encoded by this gene binds to ephrin-B2 and plays an essential role in vascular development.

Publication Note: This RefSeq record includes a subset of the publications that are available for this gene. Please see the Entrez Gene record to access additional publications.

FEATURES	Location/Qualifiers
source	1..987 /organism="Homo sapiens" /db_xref="taxon:9606" /chromosome="7" /map="7q22"
<u>Protein</u>	1..987 /product="ephrin receptor EphB4 precursor" /EC_number="2.7.10.1" /note="hepatoma transmembrane kinase; ephrin receptor EphB4"
<u>sig_peptide</u>	1..15 /calculated_mol_wt=1647
<u>mat_peptide</u>	16..987 /product="ephrin receptor EphB4" /calculated_mol_wt=106642
<u>Region</u>	29..197 /region_name="Ephrin_lbd" /note="Ephrin receptor ligand binding domain. The Eph receptors, which bind to ephrins pfam00812 are a large family of receptor tyrosine kinases. This family represents the amino terminal domain which binds the ephrin ligand; cl02704" /db_xref="CDD:93072"
<u>Region</u>	258..299 /region_name="GCC2_GCC3" /note="GCC2 and GCC3; pfam07699" /db_xref="CDD:87338"
<u>Region</u>	324..429 /region_name="FN3" /note="Fibronectin type 3 domain; One of three types of internal repeats found in the plasma protein fibronectin. Its tenth fibronectin type III repeat contains an RGD cell recognition sequence in a flexible loop between 2 strands. Approximately 2% of all...; cd00063" /db_xref="CDD:28945"
<u>Site</u>	order(324,398,413) /site_type="other" /note="Interdomain contacts" /db_xref="CDD:28945"
<u>Site</u>	order(417..418,421..422) /site_type="other" /note="Cytokine receptor motif" /db_xref="CDD:28945"
<u>Region</u>	434..526 /region_name="FN3" /note="Fibronectin type 3 domain; One of three types of internal repeats found in the plasma protein fibronectin. Its tenth fibronectin type III repeat contains an RGD cell recognition sequence in a flexible loop between 2 strands. Approximately 2% of all...; cd00063" /db_xref="CDD:28945"

Site order(434,499,514)
 /site_type="other"
 /note="Interdomain contacts"
 /db_xref="CDD:28945"
Site order(515..516,518..519)
 /site_type="other"
 /note="Cytokine receptor motif"
 /db_xref="CDD:28945"
Region 610..878
 /region_name="PTKc_EphR_B"
 /note="PTKc_EphR_B: Protein Tyrosine Kinase (PTK) family; Ephrin Receptor (EphR) subfamily; class EphB receptors; catalytic (c) domain. The PTKc family is part of a larger superfamily that includes the catalytic domains of other kinases such as protein serine...; cd05065"
 /db_xref="CDD:88348"
Region 615..874
 /region_name="Pkinase_Tyr"
 /note="Protein tyrosine kinase; pfam07714"
 /db_xref="CDD:90343"
Site order(621..625,629,645,647,693..694,696,700,740,744..745,747,758,779..783,792,826)
 /site_type="active"
 /db_xref="CDD:88348"
Site order(621..623,625,629,645,647,693..694,696,700,745,747,758)
 /site_type="other"
 /note="ATP binding site"
 /db_xref="CDD:88348"
Site order(740,744,779..783,792,826)
 /site_type="other"
 /note="substrate binding site"
 /db_xref="CDD:88348"
Site 757..781
 /site_type="other"
 /note="activation loop (A-loop)"
 /db_xref="CDD:88348"
Region 911..969
 /region_name="SAM"
 /note="Sterile alpha motif.; Widespread domain in signalling and nuclear proteins. In EPH-related tyrosine kinases, appears to mediate cell-cell initiated signal transduction via the binding of SH2-containing proteins to a conserved tyrosine that is...; cd00166"
 /db_xref="CDD:29106"
Site 924
 /site_type="phosphorylation"
 /db_xref="CDD:29106"
Site order(943,969)
 /site_type="other"
 /note="oligomerisation interface"
 /db_xref="CDD:29106"
CDS 1..987
 /gene="EPHB4"
 /coded_by="NM_004444.4:492..3455"
 /db_xref="CCDS:CCDS5706.1"
 /db_xref="GeneID:2050"
 /db_xref="HGNC:3395"
 /db_xref="HPRD:02481"
 /db_xref="MIM:600011"

ORIGIN

```
1 melrvllcwa slaaaaleetl lntkletadl kwvtfpqvdg qweelsglde eqhsvrtyev
61 cdvqrapgqa hwlrtgwvpr rgavhvayatl rftmleclsl pragrsket ftvfyyesda
121 dtataltpaw menpyikvdt vaaehltrkr pgaeatgkvn vktlrlgpls kagfylafqd
181 qgacmallsl hlffykcagl tvnltrfpet vprelvvpva gscvvdavpa pgpspslycr
241 edgqwaeqpvg tgcscapgfe aaegntkcra caqgtfkpls gegscqpcpa nshsntigsa
301 vcqcrvgyfr artdprgpc ttpsaprsv vsrlngsslh lewsaplesg gredltyalr
361 crecrpggsc apcggdltfd pgprdlvepw vvvrqlrpdf tytfvtafn gvsslatgpv
421 pfepvnvttt revppavsdv rvtrsspssl slawavprap sgavldyevk yhekgaegps
481 svrflktsen raelrglkr asylvqvrar seagygpfgq ehhsqtqlde segwreqlal
541 iagtavvgvv lvlvvivvav lclrkqsngr eaeySDKHGQ ylighgtkvy idpftyedpn
601 eavrefakei dvsyvkieev igagefgevc rgrlkapgk escvaiktlk ggyterqrre
661 flseasingq fehpnirle gvvtnsmpvm iltefmenga ldsflrlndg qftviqlvgm
721 lrgiasgmry laemsvhrd laarnilvns nlvckvsdfg lsrfleenss dptytsslgg
781 kipirwtape aiafrkftsa sdawsygvm wevmsfgerp ywdmsnqdv naieqdyrlp
841 pppdcptslh qlmldcwqkd rnarprfpqv vsalDKMIRN paslkivare nggashplld
901 qrqphysafg svgewlraik mgryeesfaa agfgsfelvs qisaedllri gvtlaghqkk
961 ilasvqhmks qakpgtpggt ggpapqy
```

//

[Disclaimer](#) | [Write to the Help Desk](#)
[NCBI](#) | [NLM](#) | [NIH](#)

Last update: Tue, 29 Apr 2008 Rev. 126150